



spinocerebellar ataxia type 1

Spinocerebellar ataxia type 1 (SCA1) is a condition characterized by progressive problems with movement. People with this condition initially experience problems with coordination and balance (ataxia). Other signs and symptoms of SCA1 include speech and swallowing difficulties, muscle stiffness (spasticity), and weakness in the muscles that control eye movement (ophthalmoplegia). Eye muscle weakness leads to rapid, involuntary eye movements (nystagmus). Individuals with SCA1 may have difficulty processing, learning, and remembering information (cognitive impairment).

Over time, individuals with SCA1 may develop numbness, tingling, or pain in the arms and legs (sensory neuropathy); uncontrolled muscle tensing (dystonia); muscle wasting (atrophy); and muscle twitches (fasciculations). Rarely, rigidity, tremors, and involuntary jerking movements (chorea) have been reported in people who have been affected for many years.

Signs and symptoms of the disorder typically begin in early adulthood but can appear anytime from childhood to late adulthood. People with SCA1 typically survive 10 to 20 years after symptoms first appear.

Frequency

SCA1 affects 1 to 2 per 100,000 people worldwide.

Genetic Changes

Mutations in the *ATXN1* gene cause SCA1. The *ATXN1* gene provides instructions for making a protein called ataxin-1. This protein is found throughout the body, but its function is unknown. Within cells, ataxin-1 is located in the nucleus. Researchers believe that ataxin-1 may be involved in regulating various aspects of producing proteins, including the first stage of protein production (transcription) and processing RNA, a chemical cousin of DNA.

The *ATXN1* gene mutations that cause SCA1 involve a DNA segment known as a CAG trinucleotide repeat. This segment is made up of a series of three DNA building blocks (cytosine, adenine, and guanine) that appear multiple times in a row. Normally, the CAG segment is repeated 4 to 39 times within the gene. In people with SCA1, the CAG segment is repeated 40 to more than 80 times. People with 40 to 50 repeats tend to first experience signs and symptoms of SCA1 in mid-adulthood, while people with more than 70 repeats usually have signs and symptoms by their teens.

An increase in the length of the CAG segment leads to the production of an abnormally long version of the ataxin-1 protein that folds into the wrong 3-dimensional shape. This

abnormal protein clusters with other proteins to form clumps (aggregates) within the nucleus of the cells. These aggregates prevent the ataxin-1 protein from functioning normally, which damages cells and leads to cell death. For reasons that are unclear, aggregates of ataxin-1 are found only in the brain and spinal cord (central nervous system). Cells within the cerebellum, which is the part of the brain that coordinates movement, are particularly sensitive to changes in ataxin-1 shape and function. Over time, the loss of the cells of the cerebellum causes the movement problems characteristic of SCA1.

Inheritance Pattern

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. An affected person usually inherits the altered gene from one affected parent. However, some people with SCA1 do not have a parent with the disorder.

As the altered *ATXN1* gene is passed down from one generation to the next, the length of the CAG trinucleotide repeat often increases. A larger number of repeats is usually associated with an earlier onset of signs and symptoms. This phenomenon is called anticipation. Anticipation tends to be more prominent when the *ATXN1* gene is inherited from a person's father (paternal inheritance) than when it is inherited from a person's mother (maternal inheritance).

Individuals who have around 35 CAG repeats in the *ATXN1* gene do not develop SCA1, but they are at risk of having children who will develop the disorder. As the gene is passed from parent to child, the size of the CAG trinucleotide repeat may lengthen into the range associated with SCA1 (40 repeats or more).

Other Names for This Condition

- olivopontocerebellar atrophy I
- SCA1
- spinocerebellar atrophy I
- type 1 spinocerebellar ataxia

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Spinocerebellar ataxia 1
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0752120/>

Other Diagnosis and Management Resources

- GeneReview: Spinocerebellar Ataxia Type 1
<https://www.ncbi.nlm.nih.gov/books/NBK1184>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Movement--Uncoordinated
<https://medlineplus.gov/ency/article/003198.htm>
- Health Topic: Balance Problems
<https://medlineplus.gov/balanceproblems.html>
- Health Topic: Cerebellar Disorders
<https://medlineplus.gov/cerebellardisorders.html>
- Health Topic: Movement Disorders
<https://medlineplus.gov/movementdisorders.html>

Genetic and Rare Diseases Information Center

- Spinocerebellar ataxia 1
<https://rarediseases.info.nih.gov/diseases/4071/spinocerebellar-ataxia-1>

Additional NIH Resources

- National Institute of Neurological Disorders and Stroke: Ataxias and Cerebellar or Spinocerebellar Degeneration Information Page
<https://www.ninds.nih.gov/Disorders/All-Disorders/Ataxias-and-Cerebellar-or-Spinocerebellar-Degeneration-Information-Page>

Educational Resources

- Boston Children's Hospital: Dysphagia
<http://www.childrenshospital.org/conditions-and-treatments/conditions/dysphagia>
- Disease InfoSearch: Spinocerebellar ataxia 1
<http://www.diseaseinfosearch.org/Spinocerebellar+ataxia+1/6743>

- Johns Hopkins Medicine Department of Neurology and Neurosurgery: What is Ataxia?
http://www.hopkinsmedicine.org/neurology_neurosurgery/centers_clinics/movement_disorders/ataxia/conditions/
- Merck Manual Consumer Version: Coordination Disorders
<http://www.merckmanuals.com/home/brain-spinal-cord-and-nerve-disorders/movement-disorders/coordination-disorders>
- National Ataxia Foundation: Spinocerebellar Ataxia Type 1
<http://www.ataxia.org/pdf/NAF%20Web%20Content%20Publication%20SCA1.pdf>
- Orphanet: Spinocerebellar ataxia type 1
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=98755
- Washington University, St. Louis: Neuromuscular Disease Center
<http://neuromuscular.wustl.edu/ataxia/domatax.html#sca1>

Patient Support and Advocacy Resources

- Family Caregiver Alliance
<https://www.caregiver.org/>
- National Ataxia Foundation
<http://www.ataxia.org/>
- National Organization for Rare Disorders (NORD): Autosomal Dominant Hereditary Ataxia
<https://rarediseases.org/rare-diseases/autosomal-dominant-hereditary-ataxia/>
- University of Kansas Medical Center Resource List: Ataxia
<http://www.kumc.edu/gec/support/ataxia.html>

GeneReviews

- Spinocerebellar Ataxia Type 1
<https://www.ncbi.nlm.nih.gov/books/NBK1184>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?cond=%22spinocerebellar+ataxia+type+1%22+OR+%22Spinocerebellar+Ataxia%22>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28spinocerebellar+ataxia+type+1%5BTIAB%5D%29+OR+%28sca1%5BTIAB%5D%29%29+AND+%28Spino+cerebellar+Ataxias%5BMAJR%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

OMIM

- SPINOCEREBELLAR ATAXIA 1
<http://omim.org/entry/164400>

Sources for This Summary

- Globas C, du Montcel ST, Baliko L, Boesch S, Depondt C, DiDonato S, Durr A, Filla A, Klockgether T, Mariotti C, Melegh B, Rakowicz M, Ribai P, Rola R, Schmitz-Hubsch T, Szymanski S, Timmann D, Van de Warrenburg BP, Bauer P, Schols L. Early symptoms in spinocerebellar ataxia type 1, 2, 3, and 6. *Mov Disord*. 2008 Nov 15;23(15):2232-8. doi: 10.1002/mds.22288.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18759344>
- Kang S, Hong S. Molecular pathogenesis of spinocerebellar ataxia type 1 disease. *Mol Cells*. 2009 Jun 30;27(6):621-7. doi: 10.1007/s10059-009-0095-y. Epub 2009 Jun 22. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19572115>
- Matilla-Dueñas A, Goold R, Giunti P. Clinical, genetic, molecular, and pathophysiological insights into spinocerebellar ataxia type 1. *Cerebellum*. 2008;7(2):106-14. doi: 10.1007/s12311-008-0009-0. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18418661>
- Zoghbi HY, Orr HT. Pathogenic mechanisms of a polyglutamine-mediated neurodegenerative disease, spinocerebellar ataxia type 1. *J Biol Chem*. 2009 Mar 20;284(12):7425-9. doi: 10.1074/jbc.R800041200. Epub 2008 Oct 28. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18957430>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2658037/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/spinocerebellar-ataxia-type-1>

Reviewed: February 2011

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services